CONTENTS

(Abstracts/contents lists published in: Chem. Abstr.; Curr. Contents, Phys. Chem. Earth Sci., Life Sci.)

Dedication: Professor Leslie Hough xii

Die Kristall- und Molekülstrukturen der aus D-Galactose und D-Glucose durch Fischer–Sowden-Reaktion erhältlichen vier isomeren Nitroheptitole
P. Köll (Oldenburg, B.R.D.), B. Malzahn, and J. Kopf (Hamburg, B.R.D.) 1

Computer-assisted interpretation of ^{1}H-n.m.r. spectra in the analysis of the structure of oligosaccharides
E. F. Hounsell and D. J. Wright (Harrow, Gt. Britain) 19

Synthese einer Kdo-haltigen Pentasaccharidesequenz der inneren Core- und Lipid A-Region von Lipopolysacchariden
H. Paulsen and C. Krogmann (Hamburg, B.R.D.) 31

Synthesis of sweet-tasting methylthio derivatives of D-fructose and sucrose
L.-X. Gan, R. L. Whistler, and J. R. Daniel (West Lafayette, IN, U.S.A.) 45

Sucrose octabenzoate: assignment of ^{13}C and ^{1}H resonances of the sucrose moiety and the ^{13}C resonances of the carbonyl carbons. Use of ^{13}C-n.m.r. spectroscopy for the study of selective deacylation
I. J. Colquhoun, A. H. Haines, P. A. Konowicz (Norwich, Gt. Britain), and H. F. Jones (Reading, Gt. Britain) 53

Syntheses of four fatty acid esters of sucrose found in type B trichomes of *Solanum berthaultii* Hawkes (wild potato), including the major component, 6-*O*-decanoyl-3,4-di-*O*-isobutyrylsucrose
S. Oscarson and H. Ritzén (Stockholm, Sweden) 61

Iodoalkoxylation of 1,5-anhydro-2-deoxyhex-1-enitols (glycals)
D. Horton, W. Priebe, and M. Sznaidman (Columbus, OH, U.S.A.) 71

Synthesis of 2′,3′-dideoxy-2′,3′-α-methanocytidine
A. R. Beard, P. I. Butler, J. Mann, and N. K. Partlett (Reading, Gt. Britain) 87

Photolabile, spacer-modified oligosaccharides for probing malto-oligosaccharide binding sites in proteins
J. Lehmann and L. Ziser (Freiburg, F.R.G.) 93

The synthesis and resolution of (±)-1,5,6-tri-*O*-benzyl-*myo*-inositol
T. Desai, A. Fernandez-Mayoralas, J. Gigg, R. Gigg, and S. Payne (London, Gt. Britain) 105

Synthesis of methyl 3-*O*-(α-D-glucopyranosyl)-7-*O*-(L-*glycero*-α-D-*manno*-heptopyranosyl)-L-*glycero*-α-D-*manno*-heptopyranoside
P. J. Garegg, S. Oscarson, and M. Szönyi (Stockholm, Sweden) 125

Synthesis of a tri- and a tetra-saccharide fragment of the capsular polysaccharide of type III Group B *Streptococcus*
V. Pozsgay, J.-R. Brisson, and H. J. Jennings (Ottawa, Ont., Canada) 133

Stereoselective total synthesis of dodecagalacturonic acid, a phytoalexin elicitor of soybean
Y. Nakahara and T. Ogawa (Wako, Japan) 147

2,3′-Anhydrosucrose
K. Čapek, E. Čadová, and P. Sedmera (Prague, Czechoslovakia) 161

The formation of isopropylidene acetals of erythritol and ribitol under conditions of kinetic control
A. Awal, A. S. F. Boyd, J. G. Buchanan, and A. R. Edgar (Edinburgh, Gt. Britain) . . . 173

A comparison of the results of sequential hydrazinolysis–nitrosation and alkali-mediated cleavage–nitrosation of the *O*-linked oligosaccharides of gastric mucus glycoproteins
S. R. Carter, J. M. Williams (Swansea, Gt. Britain), and J. R. Clamp (Bristol, Gt. Britain) 181

Acyloxonium ions in the high-yielding synthesis of oxolanes from alditols, hexoses, and hexonolactones catalysed by carboxylic acids in anhydrous hydrogen fluoride
J. Defaye, A. Gadelle (Grenoble, France), and C. Pedersen (Lyngby, Denmark) 191

Synthesis and reactions of chlorodeoxy-L-talofuranoid derivatives
C.-K. Lee (Kent Ridge, Singapore) 203

Halogenation reactions of derivatives of D-glucose and sucrose
R. Khan and G. Patel (Reading, Gt. Britain) 211

Synthesis of luciferin glycosides as substrates for novel ultrasensitive enzyme assays
R. Amess, N. Baggett, P. R. Darby, A. R. Goode (Birmingham, Gt. Britain), and E. E. Vickers (Suffolk, Gt. Britain) 225

Synthesis of oligosaccharins: a chemical synthesis of propyl *O*-β-D-galactopyranosyl-(1→2)-*O*-α-D-xylopyranosyl-(1→6)-*O*-β-D-glucopyranosyl-(1→4)-β-D-glucopyranoside
A. Wotovic, J.-C. Jacquinet (Orleans, France), and P. Sinaÿ (Paris, France) 235

Synthetically useful transformations of a branched-chain, deoxynitro-1-thio-D-glucitol, including a novel mode of 1-thioglycoside formation
H. H. Baer and F. Santoyo González (Ottawa, Ont., Canada) 247

Synthesis of some monodeoxy- and dideoxy-hexitols, and derivatives thereof, from D-glucono-1,5-lactone
H. Regeling and G. J. F. Chittenden (Nijmegen, The Netherlands) 261

Synthesis of the enantiomers of 6-epicastanospermine and 1,6-diepicastanospermine from D- and L-gulonolactone
G. W. J. Fleet, N. G. Ramsden (Oxford, Gt. Britain), R. J. Nash, L. E. Fellows (Surrey, Gt. Britain), G. S. Jacob (St. Louis, MO, U.S.A.), R. J. Molyneux (Albany, CA, U.S.A.), I. Cenci di Bello, and B. Winchester (London, Gt. Britain) 269

Hexa-*O*-benzyl-5-hydroxy-pseudo-α-D-glucopyranose and its C-5 epimer
R. J. Ferrier and A. E. Stüz (Wellington, New Zealand) 283

An enantiospecific synthesis of (−)-talaromycins A and B from D-fructose
I. Izquierdo Cubero and M. T. Plaza López-Espinosa (Granada, Spain) 293

Formation of disaccharides related to heparin and heparan sulfate by chemial modification of maltose
J. N. Glushka and A. S. Perlin (Montreal, Que., Canada) 305

Oxidation periodique de derivés du saccharose
A. Badel, G. Descotes, and J. Mentech (Villeurbanne, France) 323

Synthesis and reactions of leucrose and its exocyclic glycal
J. Thiem and M. Kleeberg (Münster, F.R.G.) 333

Structure of the amino acid-containing capsular polysaccharide from *Escherichia coli* O8:K49:H21
L. M. Beynon, G. G. S. Dutton (Vancouver, B.C., Canada), and J. C. Richards (Ottawa, Ont., Canada) 347

The structure of the capsular antigen from *Escherichia coli* O8:K87:H19
H. Parolis, L. A. S. Parolis, S. M. R. Stanley (Grahamstown, South Africa), and G. G. S. Dutton (Vancouver, B.C., Canada) 361

Identification of the lipopolysaccharide O-chain of *Escherichia hermannii* (ATCC 33651) as a D-rhamnan
M. B. Perry and J. C. Richards (Ottawa, Ont., Canada) 371

Notes

Structural and conformational studies of 2,7-anhydro-L-*glycero*-β-D-*manno*-octulopyranose and its per-*O*-acetyl derivative
M. G. Ambrose, A. J. Fatiadi, and B. Coxon (Gaithersburg, MD, U.S.A.) 377

Synthesis of 4-nitrophenyl *O*-(2-acetamido-2-deoxy-β-D-glucopyranosyl)-(1→2)-*O*-(4-*O*-methyl-α-D-mannopyranosyl)-(1→6)-β-D-glucopyranoside. A potential specific acceptor-substrate for *N*-acetylglucosaminyltransferase-V (GnT V)
S. H. Khan, S. A. Abbas, and K. L. Matta (Buffalo, NY, U.S.A.) 385

A novel synthesis of methyl 2-*O*-methyl-α,β-L-fucofuranoside
M. K. Gurjar and N. Ramaiah Srinivas (Hyderabad, India) 398

N.m.r. method for the identification of sucrose acetates
E. B. Rathbone (Reading, Gt. Britain) 402

Selective deacetylation of sucrose octa-acetate with primary amines to give 2,3,4,6,1′,6′-hexa-*O*-acetylsucrose
A. H. Haines, P. A. Konowicz (Norwich, Gt. Britain), and H. F. Jones (Reading, Gt. Britain) 406

New observations on the Garegg–Samuelsson olefination reaction
Z. Pakulski and A. Zamojski (Warsaw, Poland) 410

Synthesis and resolution of 2,3-diamino-1-cyclohexanols
K. Izawa, T. Ineyama (Kawasaki, Japan), K. Fujii, and T. Suami (Yokohama, Japan) . . 415

Double asymmetric induction in the catalytic osmylation of some α,β-unsaturated octuronic acid derivatives
J. S. Brimacombe, G. McDonald, and M. Abdur Rahman (Dundee, Gt. Britain) 422

Titanium-mediated methylene transfer reactions on sugar esters, lactones, and uloses
M. H. Ali, P. M. Collins, and W. G. Overend (London, Gt. Britain) 428

Synthesis of phenyl 4-(*N*-benzylformamido)-4,6-dideoxy-3-*C*-methyl-2-*O*-methyl-1-thio-α-L-mannopyranoside, an *N*-formylkansosamine glycosyl donor
I. Bajza and A. Lipták (Debrecen, Hungary) 435

Structural studies of the O-antigen polysaccharide of *Vibrio fluvialis* AA-18239
L. Kenne, B. Lindberg, M. Mahbubur Rahman (Stockholm, Sweden), and M. Mosihuzzaman (Dhaka, Bangladesh) 440

Studies on cervical glycoproteins. Isolation and characterization of neutral oligosaccharides from Pronase-treated glycoproteins of bonnet monkey (*Macacca radiata*)
Nasir-ud-Din, S. Altaf Hussain (Quetta, Pakistan), R. W. Jeanloz (Boston, MA, U.S.A.), and E. Walker-Nasir (Geneva, Switzerland) 444

Preliminary communications

Use of n.m.r. imaging to map the spatial distribution of structure in polysaccharide gels
S. L. Duce, T. A. Carpenter, and L. D. Hall (Cambridge, Gt. Britain) c1

New colorimetric substrates for the assay of glycosidases
K. H. Aamlid, G. Lee, B. V. Smith, A. C. Richardson, and R. G. Price (London, Gt. Britain) c5

Regioselective hydrogenolysis of benzyl glycosides
T. Bieg and W. Szeja (Gliwice, Poland) c10

The effect of substituting key hydroxyl groups by amino groups on the binding of the Lewis b tetrasaccharide by a lectin and a monoclonal antibody
R. U. Lemieux, R. Szweda, E. Paszkiewicz-Hnatiw, and V. Spohr, (Edmonton, Alb, Canada) c12

Author index c19

Subject index c21